Searches for new interstellar molecules, including a tentative detection of aziridine and a possible detection of propenal

著者	DICKENS J.E, IRVINE W.M, NUMMELIN A, MOLLENDAL H, SAITO S, THORWIRTH S, HJALMARSON A, OHISHI M
journal or publication title	Spectrochimica Acta - A - Molecular and Biomolecular Spectroscopy
volume	57
number	4
page range	643-660
year	2001-03
URL	http://hdl.handle.net/10098/1574

doi: 10.1016/S0029-5493(01)00333-8

Searches for new interstellar molecules, including a tentative detection of aziridine and a possible detection of propenal

J.E. Dickens ^a, W.M. Irvine ^b, A. Nummelin ^c, H. Møllendal ^d, S. Saito ^e,
S. Thorwirth ^f, Å Hjalmarson ^g, M. Ohishi ^h

^a Jet Propulsion Lab., MS 169-506, 4800 Oak Grove Dr., Pasadena, CA 91109, USA

^b Department of Astronomy, 619 LGRT, University of Massachusetts, Amherst, MA 01003, USA

^c Department of Physics, Rensselaer Polytechnic Institute, Troy, NY 12180-3590, USA

^d Department of Chemistry, University of Oslo, PO Box 1033, Blindern, N-0315 Oslo, Norway

^e Research Center for Development of Far-IR Region, Fukui University, 3-9-1 Bunkyo, Fukui 910-8507, Japan

^f I. Physikalisches Institut, Universität zu Köln, D50937 Köln, Germany

^g Onsala Space Observatory, Chalmers University of Technology, S-43992 Onsala, Sweden

^h National Astronomical Observatory of Japan, 2-21-1, Ohsawa, Mitaka, Tokyo, 181, Japan

Abstract

Rotational spectroscopy at millimeter wavelengths is a powerful means of investigating the chemistry of dense interstellar clouds. These regions can exhibit an interesting complement of gas phase molecules, including relatively complex organics. Here we report the tentative first astronomical detection of aziridine (ethylenimine), the possible detection of propenal (acrolein), and upper limits on the abundances of cyclopropanone, furan, hydroxyethanal (glycolaldehyde), thiohydroxylamine (NH₂SH), and ethenol (vinyl alcohol) in various interstellar clouds.

Keywords: Interstellar molecules; Rotational spectra; Propenal; Aziridine; Cyclopropanone; Furan; Hydroxyethanal

1. Introduction

Over the last three decades astronomers have discovered that the denser clouds in the interstellar medium (ISM) have a complex chemistry involving both particulate matter (interstellar

grains) and a gas phase in which more than 100 molecular species have been securely identified [1–3]. The latter include molecules with up to at least 13 atoms and molecular weights up to 147 (Table 1). In addition, there is strong evidence for even heavier aromatic molecules, although specific assignments have been difficult to achieve [4].

This interstellar chemistry is scientifically important, both in its own right as it pertains to the largest objects in the universe whose composition

is subject to chemical bonding; and in its application to astrophysical problems such as star formation, where cooling of clouds by trace molecular constituents and the coupling to magnetic fields through the abundance of ions are critically important [5]. Moreover, the low temperatures (often ≈ 10 K) and densities ($\geq 10^4$ molecules cm^{-3} in 'dense' regions) make these clouds interesting laboratories where the spectra and structure of a number of molecular species have been studied before this was possible in the laboratory (e.g. [6–8]). Finally, since interstellar organic molecules may well be incorporated into comets and hence brought to the early Earth, it is at least conceivable that such exogenous ma-

terial may have played some role in the origin of life [15,53].

Although much is known about the chemistry of dense interstellar clouds, much also remains uncertain, as a result of the variety of heterogeneous processes which may be important, including the effects of radiation fields and cosmic rays. Absolute abundances are difficult to determine as a function of physical conditions; such conditions vary significantly among and within different environments; reaction rates and branching ratios in the gas phase may not be known at relevant temperatures; the nature of grain surfaces and hence processes on and in grains are poorly constrained (although spectra

Table 1
Gas phase interstellar and circumstellar molecules

2 Atoms	3 Atoms	4 Atoms	5 Atoms	6 Atoms	7 Atoms	8 Atoms	9 Atoms	≥ 10 Atoms
H ₂	C ₂ H	C ₂ H ₂	C ₄ H	C ₂ H ₄ ^a	C ₆ H	CH ₃ COOH	CH ₃ C ₄ H	CH ₃ C ₄ CN ^b
C ₂	CH ₂	l-C ₃ H	C ₃ H ₂	H ₂ CCCC	HC ₅ N	HCOOCH ₃	CH ₃ OCH ₃	CH ₃ COCH ₃
CH	HCN	c-C ₃ H ^c	H ₂ CCC	CH ₃ OH	CH ₂ CHCN	C ₇ H ^a	CH ₃ CH ₂ CN	HC ₉ N
CH ⁺	HNC	NH ₃	HCOOH	CH ₃ CN	CH ₃ C ₂ H	H ₂ C ₆	CH ₃ CH ₂ OH	HC ₁₁ N
CN	HCO	HNCO	CH ₂ CO	CH ₃ NC	CH ₃ CHO	HC ₆ H ^a	HC ₇ N	(C ₂ H ₅) ₂ O ^b
CO	HCO ⁺	HOCO ⁺	HC ₃ N	CH ₃ SH	CH ₃ NH ₂	CH ₃ C ₃ N	C ₈ H	
CS	HOC ⁺	HCNH ⁺	CH ₂ CN	NH ₂ CHO	c-CH ₂ OCH ₂ ^c			
OH	N ₂ H ⁺	HNCS	NH ₂ CN	HC ₃ HO				
NH	NH ₂	C ₃ N	CH ₂ NH	C ₅ H				
NO	H ₂ O	C ₃ O	CH ₄	HC ₃ NH ⁺				
NS	HCS ⁺	H ₂ CS	SiH ₄ ^a	C ₅ N				
SiC ^a	H ₂ S	C ₃ S	C ₄ Si ^a	HC ₄ H ^a				
SiO	OCS	HCCN	C ₅ ^a					
SiS	N ₂ O	H ₃ O ⁺	HCCNC					
SiN ^a	SO ₂	H ₂ CN	HNCCC					
SO	SiC ₂ ^a	H ₂ CO	H ₂ COH ⁺					
HCl	C ₂ S	c-SiC ₃ ^c						
CP ^a	C ₂ O	CH ₃						
SO ⁺	C ₃							
NaCl ^a	MgNC ^a							
AlCl ^a	MgCN ^a							
KCl ^a	NaCN ^a							
AlF ^a	HNO							
PN	H ₃ ⁺							
CO ⁺								
SiH ^b								
HF ^b								

^a Detected only in stellar envelopes.

^b Tentative detection.

^c C-, cyclic structure. For references, see [50,61,62].

Table 2
Source information^a

Source	$\alpha(1950.0)$	$\delta(1950.0)$	V_{LSR} [km s ⁻¹] ^b	$N(\text{H}_2)$ [cm ⁻²] ^c	θ_{mb} [arcsec] ^d	Reference/comments
TMC-1(CP)	04 ^h 38 ^m 38.6 ^s	+25°35'45.0''	+5.8	2.0(22)	50	[23] from C ¹⁸ O
Orion KL	05 ^h 32 ^m 47.0 ^s	-05°24'23.0''	+9.0	1.0(23)	50	[28] from C ¹⁸ O
Orion 3'N	05 ^h 32 ^m 51.0 ^s	-05°20'50.0''	+9.0	5.0(22)	50	[28] from C ¹⁸ O
G327.3-0.6	15 ^h 49 ^m 15.6 ^s	-54°28'07.0''	-45.0	3.0(23)	45	[51] from C ¹⁸ O
L134N(C)	15 ^h 51 ^m 32.6 ^s	-02°43'31.0''	+2.5	1.5(22)	50	[24] from C ¹⁸ O
IRS 16293	16 ^h 29 ^m 20.9 ^s	-24°22'13.0''	+4.0	2.0(23)	20	[25] from C ¹⁷ O
NGC 6334F	17 ^h 17 ^m 32.3 ^s	-35°44'02.5''	-7.0	2.0(23)	45	[21] from C ¹⁷ O
SgrB2 N	17 ^h 44 ^m 10.1 ^s	-28°21'17.0''	+68.0	3.0(24)	23	[18] from C ¹⁷ O
G10.47+0.03	18 ^h 05 ^m 40.3 ^s	-19°52'21.0''	+68.0	1.0(23)	45	[21] from C ¹⁷ O
W33A	18 ^h 14 ^m 26.3 ^s	-17°52'01.0''	+32.0	6.2(23)	14	[52] from C ¹⁷ O
G31.41+0.31	18 ^h 44 ^m 59.2 ^s	-01°16'07.0''	+97.0	2.0(23)	45	[21] from C ¹⁷ O
G34.3+0.2	18 ^h 50 ^m 46.2 ^s	+01°11'13.0''	+58.0	3.0(23)	45	[21] from C ¹⁷ O

^a Note: notation a(b) represents a $\times 10^b$.

^b V_{LSR} is the radial velocity of the source with respect to the local standard of rest (see text).

^c $N(\text{H}_2)$ is the column density of molecular hydrogen along the line of sight as estimated from the CO isotopomer indicated in the last column.

^d θ_{mb} is the antenna main beam size (full width half power).

show that the grains contain both silicates and poorly characterized organic material [5,74]; and the interchange of material between solid and gas phases is not well understood [9,10]. Thus, the additional constraints on models supplied by the identification of new molecular species and the determination of their abundance remains very important.

The increased sensitivity of infrared detectors and the necessity of such measurements for non-polar species and for solid state features are increasing the importance of infrared observations of vibrational spectra for interstellar chemistry (e.g. [11,12]). Nonetheless, the great bulk of the relevant data come from heterodyne observations of rotational spectra at millimeter and sub-millimeter wavelengths. Rotational spectroscopy is thus critical to this field, and key data are often lacking, particularly for 'unstable' species such as radicals and positive ions, or even for well known molecules in high rotational levels, in low-lying vibrational states, and for rarer isotopomers (cf. the recent important work on species such as methanol and methyl formate (methyl methanate; e.g. [13,14]).

Here we report on searches for several previously undetected molecules in a number of interstellar sources. These particular searches were

motivated by the identification of related species in interstellar clouds or, in one case, in a comet (the chemistry of cometary comae is remarkably similar to that in the ISM, either because of the presence of preserved interstellar material or because of similar conditions in the ISM to those in the region of cometary formation in the solar nebula [15]). We have taken advantage of the relatively large body of data which we accumulated in several searches targeted for certain molecules, to seek for other species with transitions in the frequency ranges encompassed by the data. Our observed frequency bands fall within the general ranges 18–23, 85–115, 130–170, and 210–260 GHz (wavelengths of about 17–13, 3.5–2.6, 2.3–1.8, and 1.4–1.2 mm, respectively).

Observations were made at the 15 m diameter Swedish-ESO Submillimeter Telescope (SEST) in Chile, the US National Radio Astronomy Observatory's 12 m telescope in Arizona, and the NASA DSN 70 m telescope. Our observed sources (Table 2) are primarily examples of the regions called hot cores, which are present around very young, massive stars or protostars (e.g. [2,16,18]). These hot cores include Orion KL (the closest such region to the Sun, and

2. Molecular species sought

We give our results consecutively for each molecular species. Table 3 lists the frequencies, energies above the ground state, and Einstein A-coefficients for the transitions observed for each species except aziridine, for which the transitions are given in Table 4. Because traditionally radio astronomers worked at frequencies ν for which the Rayleigh–Jeans approximation was valid ($h\nu \ll kT$), it is usual to express intensity (I) in terms of temperature units via

$$I(\nu) = (2h\nu^3/c^2)[\exp(h\nu/kT) - 1]^{-1} \approx 2k\nu^2 T/c^2,$$

where the ‘brightness temperature’ T is the temperature of a hypothetical blackbody filling the beam of a perfect antenna and producing the measured monochromatic flux. Since a real antenna is not perfect and various calibration schemes are used, the observed signal defines an ‘antenna temperature’ T_a^* (e.g. Fig. 1) which is related to the brightness temperature by an efficiency factor. A particular choice of efficiency factor defines the ‘main beam antenna temperature’ T_{mb} (Tables 4–6, Figs. 2–5) [9,31].

Table 4 and Table 5 list for aziridine and propenal, respectively, the detected emission lines or the upper limits in each observed source region in units of milliKelvin. Table 6 lists for thiohydroxylamine the upper limits only on signal strength in each observed source region. The upper limits on line intensity for the other species which we have sought are comparable to those for thiohydroxylamine and are not tabulated; instead we give below limits on the derived column densities and abundances. Of particular interest, we report the tentative first astronomical detection of aziridine and a possible first astronomical detection of propenal.

2.1. Aziridine (ethylenimine, $c\text{-C}_2\text{H}_4\text{NH}$)

Cyclic molecules play a fundamental role in biochemistry. In the diffuse interstellar medium there is strong evidence for the presence of polycyclic aromatic hydrocarbons (PAHs), but specific molecular species have not been identified [4]. In dense interstellar clouds, cyclic species have been

hard to find; until recently only the 3-carbon rings C_3H and C_3H_2 had been identified, although two small silicon-containing rings are known in the envelopes of evolved, carbon-rich stars (Table 1). The recent identification of oxirane (ethylene oxide, $c\text{-C}_2\text{H}_4\text{O}$) in the ISM [21,29] made it natural to investigate the possible presence of the related species aziridine, in which the oxygen is replaced by an NH group. Fortunately, the rotational spectrum at relevant wavelengths has recently been measured [33,63]. Although a targeted search for aziridine was not carried out, a number of its transition frequencies were covered in our searches for other molecular species. The predicted relative intensity of these lines of course depends on the assumed rotational temperature. One of the lines which we would expect to be among the strongest at temperatures in the range 30–100 K, which is typically what we would expect in hot molecular cloud cores, is the $9_{09}\text{--}8_{18}/9_{19}\text{--}8_{08}$ doublet at 249 419.6 MHz. This line is unfortunately not observable, being less than 0.4 MHz from the $15(3)\text{--}15(2)$ A-+ line of methanol, which is a very prominent emission feature in all hot cores.

In our data there are two features that may be attributed to aziridine (Table 4). The first is the $5_{50}\text{--}4_{41}$ line at 226 233.4 MHz, where, to the best of our knowledge, no other known interstellar species have transitions. There is a clear emission line here in the spectra of the hot cores G327.3, G10.47, and Sgr B2(N), and a weak line in G34.3 (Fig. 2). The second feature is the $2_{11}\text{--}1_{01}$ transition at 86 312.7 MHz, where a line is detected in G327.3, G10.47, and possibly in G31.41 (Fig. 3). The significance of these detections will be discussed below.

2.2. Propenal (acrolein, CH_2CHCHO)

The presence in a variety of interstellar clouds of both CH_2CHCN and CH_3CHO (e.g. [21,30]) suggests that propenal might also be present. Note that ethene (H_2CCH_2) has been detected in stellar envelopes through its vibrational spectrum in absorption [61], but its non-polar nature precludes detection of rotational lines in cold interstellar clouds. The *trans*-form of propenal is more

Table 3
Molecular transition data

Rest frequency [MHz]	Transition [$J_{K-1, K+1}$]	E_u [K] ^a	A_{ul} ^b	Telescope
CH₂CHOH:ethenol				
77726.6	4 _{0,4} -3 _{0,3}	9	[10 ⁻⁵ s ⁻¹] 0.092	NRAO
84401.7	8 _{1,7} -8 _{0,8}	37	0.167	SEST
86557.6	2 _{1,2} -1 _{0,1}	5	0.074	NRAO
101449.2	5 _{1,4} -4 _{1,3}	17	0.201	SEST
222996.9	4 _{2,3} -3 _{1,2}	19	1.758	NRAO
238774.0	12 _{1,12} -11 _{0,11}	72	3.620	NRAO
240593.6	12 _{1,11} -11 _{1,01}	78	2.925	SEST
240786.7	13 _{1,13} -12 _{1,12}	84	2.948	SEST
249262.5	3 _{3,0} -3 _{2,1}	27	1.497	SEST
249560.1	4 _{3,2} -4 _{2,3}	31	2.112	SEST
249773.7	5 _{3,3} -5 _{2,4}	36	2.433	SEST
250143.9	6 _{3,4} -6 _{2,5}	41	2.632	SEST
c-C₃H₂O:cyclopropenone				
23105.6	5 _{1,4} -5 _{1,5}	12	[10 ⁻⁵ s ⁻¹] 0.009	DSN
74052.8	5 _{1,4} -4 _{1,3}	12	3.971	NRAO
82283.2	6 _{0,6} -5 _{0,5}	14	5.727	NRAO
103069.9	7 _{1,6} -6 _{1,5}	21	10.199	NRAO
105476.5	8 _{1,8} -7 _{1,7}	24	12.170	NRAO
107805.0	8 _{0,8} -7 _{0,7}	24	13.101	SEST
c-C₄H₄O:furan				
23402.5	4 _{3,2} -4 _{1,3}	8	[10 ⁻⁶ s ⁻¹] 0.012	DSN
23453.1	2 _{0,2} -1 _{0,1}	2	0.023	DSN
70074.7	6 _{1,5} -5 _{1,4}	13	0.697	NRAO
88748.7	9 _{0,9} -8 _{0,8}	22	1.786	SEST
107431.0	11 _{0,11} -10 _{0,10}	32	3.232	SEST
210175.6	22 _{0,22} -21 _{0,21}	118	25.31	NRAO
CH₂OHCHO:hydroxyethanal				
17981.1	3 _{1,2} -3 _{0,3}	4	[10 ⁻⁷ s ⁻¹] 2.132	DSN
22142.9	4 _{1,3} -4 _{0,4}	6	3.453	DSN
22252.8	9 _{3,6} -8 _{4,5}	30	0.436	DSN
23415.7	1 _{1,1} -0 _{0,0}	1	3.712	DSN
23415.9	3 _{0,3} -2 _{1,2}	2	2.080	DSN
CH₂CHCHO:propenal				
88523.3	10 _{0,10} -9 _{0,9}	23	[10 ⁻⁵ s ⁻¹] 2.753	SEST
88540.6	32 _{5,27} -33 _{4,30}	277	0.022	SEST
89043.5	10 _{6,4} -9 _{6,3}	98	1.794	SEST
89082.2	10 _{3,8} -9 _{3,7}	42	2.554	SEST
89093.2	10 _{3,7} -9 _{3,6}	42	2.555	SEST
94414.6	21 _{1,20} -21 _{1,21}	102	0.016	NRAO
97263.1	11 _{0,11} -10 _{0,10}	28	3.669	NRAO
107404.2	21 _{1,20} -21 _{0,21}	102	0.119	SEST
107528.8	12 _{2,10} -11 _{2,9}	42	4.845	SEST
NH₂SH:thiohydroxylamine				
84243.6	3 _{1,3} -2 _{1,2}	14	[10 ⁻⁶ s ⁻¹] 2.349	SEST

^a E_u is the energy of the upper level above the ground state for the transition.

^b A_{ul} is the rate of spontaneous emission from the upper to lower level, also called the Einstein A -coefficient.

stable than the *cis*-form, which lies about 700 cm^{-1} higher in energy. The rotational spectrum of *trans*-propenal is known from laboratory measurements [32].

Although we did not specifically target propenal in any of our searches, six unassigned emission lines which we observed corresponded very well in frequency to transitions of *trans*-propenal (Table 5 and Figs. 4 and 5). The significance of these results is discussed in Section 3 below.

2.3. Cyclopropenone (*c*- $\text{C}_3\text{H}_2\text{O}$)

Cyclopropenylidene (*c*- C_3H_2) is ubiquitous and relatively abundant in a variety of interstellar and circumstellar environments, and possibly in comets as well [37–39]. The linear carbon chains C_3O and C_2O are less common, but are found [40,41]. It is therefore natural to search for interstellar cyclopropenone (*c*- $\text{C}_3\text{H}_2\text{O}$), in which the unbonded electrons of *c*- C_3H_2 are shared with an oxygen atom. We used the transition frequencies measured by Guillemin et al. [42]. Several transitions in the range from 23 to 108 GHz were sought (Table 3), but only upper limits were obtained. The procedure for obtaining upper limits on the column density toward each observed source (Table 7) is described in Section 3.3 below.

2.4. Ethenol (vinyl alcohol, CH_2CHOH)

Ethenol is an isomer of two known interstellar molecules, oxirane (*c*- $\text{C}_2\text{H}_4\text{O}$) and ethanal (acetaldehyde, CH_3CHO). It has a heat of formation intermediate between those two species and hence might be thought to be intermediate in stability. Although other structural isomers are found in the ISM (e.g. HCN/HNC , $\text{CH}_3\text{CN/CH}_3\text{NC}$, cyclic- C_3H_2 /linear- H_2C_3 , $\text{CH}_2\text{CH}_2\text{OH}/(\text{CH}_3)_2\text{O}$), the detection of ethenol would provide the first triplet of such isomers. Laboratory data obtained by Saito [43] have been augmented by his more recent, higher frequency measurements for the present study. A number of transitions in the 3 mm and 1 mm bands were sought in several sources, but no lines were detected. The corresponding upper limits on column density are given in Table 7.

2.5. Furan ($\text{C}_4\text{H}_4\text{O}$)

As noted above, the only cyclic species unambiguously identified to date in interstellar clouds are the 3-membered rings C_3H_2 , C_3H , and *c*- $\text{C}_2\text{H}_4\text{O}$. As there is substantial evidence for polycyclic aromatic species from infrared spectral features that can be assigned to functional groups, however, it is natural to seek larger ring molecules in dense clouds. Searches for, e.g. furan, pyrrole

Table 4
Aziridine Observations

Transition	Frequency (MHz)	E_u (K) ^a	A_{ul} (10^{-5} s^{-1}) ^a	G327.3 T_{mb} (mK) ^a	NGC633 T_{mb} (mK) ^a	G10.47 T_{mb} (mK) ^a	G31.41 T_{mb} (mK) ^a	G34.3 T_{mb} (mK) ^a	SgrB2 T_{m} (mK) ^a
2 ₁₁ –1 ₀₁	86312.7	6	0.413	51	<10	32	17		
3 ₁₃ –2 ₀₂	88964.3	9	0.269	<4	<4				
3 ₀₃ –2 ₁₂	88739.7	9	0.269	<4	<4				
8 ₃₅ –8 ₂₆	93472.9	65	0.147	<9					
8 ₄₅ –8 ₃₆	93494.1	65	0.147	<9					
5 ₀₅ –4 ₁₄	142374.3	21	1.28	^b	^b	^b	^b		
5 ₁₅ –4 ₀₄	142375.6	21	1.28	^b	^b	^b	^b		
5 ₅₀ –4 ₄₁	226233.4	32	3.99	90	<22	79	<20	27	290

^a E_u is energy of the upper level of the transition above the ground state; A_{ul} is the Einstein A-coefficient; the last 6 columns give the observed main beam antenna temperatures (see text, [31]) or 1σ upper limits toward the 6 indicated sources (see Table 2).

^b Possible blending with the 12–11 transition of OC^{34}S makes aziridine upper limit uncertain.

Table 5

Propenal possible detections and upper limits^a

Source	$T_{\text{mb}}(\text{mK})$								
	$32_{5,27}$ – $33_{4,30}$	$10_{6,4}$ – $9_{6,3}$	$10_{3,8}$ – $9_{3,7}$	$10_{3,7}$ – $9_{3,6}$	$21_{1,20}$ – $21_{1,21}$	$21_{1,20}$ – $21_{0,21}$	$10_{0,10}$ – $9_{0,9}$	$11_{0,11}$ – $10_{0,10}$	$12_{2,10}$ – $11_{2,9}$
G327.3–0.6	15.0	9.0	13.0	23.0	30.0	18.0	<3.4	<6.4	<5.0
NGC 6334F	<3.6	<3.6	<3.6	<3.6	<7.0	<5.0	<3.6	<6.0	<4.5
SgrB2N	<9.0	44.0	51.0	99.0	...	59.0	<6.5	...	<6.0
G10.47+0.03	...	...	...	...	<7.0	...	...	<6.5	...
G31.41+0.31	...	...	...	...	13.0	...	...	<6.5	...

^a Note: Entries with < are 1σ rms upper limits.

and imidazole [44–46] had been carried out some years ago, but instrumental sensitivity has improved substantially since that time. Furan has a special significance for prebiotic chemistry, since this ring forms the basis of the simple sugars ribose and deoxyribose, the backbone molecules of RNA and DNA. Molecular constants accurate enough to predict furan transition frequencies for $J < 30$ to better than 60 kHz have been measured by Włodarczyk et al. [47]. We sought transitions over a wide range of frequencies and energies above ground (Table 3) in several astronomical sources. Only upper limits were obtained, from which the column density limits in Table 7 were derived.

2.6. Hydroxyethanal (glycolaldehyde, HOCH_2CHO)

Hydroxyethanal is an isomer of both methyl formate (methyl methanate, HCOOCH_3) and acetic acid (ethanoic acid, CH_3COOH). Methyl formate is widespread among regions of massive star formation, with a very large number of detected transitions at millimeter wavelengths (e.g. [54,17,18]). Acetic acid, in contrast, has probably been detected in only a single source [55]. Ab initio calculations were carried out for the preferred conformers of acetic acid [67], methyl formate [68] and hydroxyethanal [49] using the correlation-consistent polarized triple-zeta basis set with diffuse functions [69] (AUG-cc-pVTZ) in the Møller–Plesset (MP) second-order perturbation procedure [70] with full geometry optimization. The Gaussian 98 program [71] was employed in these computations.

Interestingly, these very high level MP2/AUG-cc-pVTZ calculations predict acetic acid (total energy: $-600\,511.17\text{ kJ mol}^{-1}$) to be about 72 kJ mol^{-1} more stable than methyl formate, showing that the formation of methyl formate is controlled by kinetic factors and not by thermodynamics (as is expected under typical conditions in interstellar clouds). Hydroxyethanal is the highest energy isomer of the 3, being about 118 kJ mol^{-1} less stable than acetic acid. Although, as noted in Section 2.4, other structural isomeric pairs are found in the ISM, no triplets have been identified. Because of the likelihood of related pathways for production and destruction of such isomers, determination of their relative abundance should place significant constraints on chemical models. Hydroxyethanal is also important as a potential intermediary in the prebiotic syntheses of sugars and sugar phosphates (e.g. [48]).

Our search was carried out exclusively at the NASA DSN 70 m telescope, at frequencies in the 18–23 GHz range (Table 3), using the transition frequencies given by Marstokk and Møllendal [49], supplemented by more recent laboratory data by one of us (HM). No emission features were detected which corresponded to transitions of hydroxyethanal. The relationship of our limits to a recent reported detection of hydroxyethanal [73] is discussed in Section 4 below.

2.7. Thiohydroxylamine (NH_2SH)

The first cometary molecule containing both nitrogen and sulfur, the NS radical, was recently

identified in the coma (atmosphere) of comet Hale–Bopp [34]. It is unclear whether such a radical could be stored in the nuclear ices over the lifetime of the solar system. It seems more likely that it is either the photodissociation product of some larger precursor, or the product of chemistry in the coma [35]. In the former case a logical precursor might be NH_2SH , since such simpler species as H_2S and NH_3 are relatively abundant in comets [15] as well as in the ISM. The microwave spectrum of NH_2SH was measured by Lovas et al. [36] and was refit by one of us (SS). We did not search specifically for NH_2SH . However, a transition expected to be strong at the temperatures of interstellar clouds was covered in our observations targeted toward other molecules. This enables us to place upper limits on the abundance of NH_2SH in several sources (Table 6).

3. Results

From the intensities of the emission lines which we have detected, or the corresponding upper limits, we now wish to determine the column density or its limit for each molecular species toward each source for which we have data.

3.1. Aziridine (ethylenimine, $c\text{-C}_2\text{H}_4\text{NH}$)

As discussed in Section 2.1, we have detected two lines in several sources which may be assignable to aziridine. Assuming a rotational temperature of 30–50 K and optically thin (unsaturated) emission, the standard relationship between observed line intensity and the corresponding column density

(e.g. [57]) shows that both of these lines are consistent with aziridine column densities of order 10^{14} cm^{-2} for the two hot cores, G327.3 and G10.47. However, there are other lines that at first glance should have been detected but are not seen, most notably the $5_{05}-4_{14}/5_{15}-4_{04}$ doublet at 142 375 MHz, but also, e.g. the $3_{03}-2_{12}$ and $3_{13}-2_{02}$ lines at 88 964.3 and 88 739.7 MHz, respectively (Table 4). The lack of emission features at these frequencies might imply an aziridine abundance 2–5 times lower than the one inferred from the two detected lines, although part of the absence of the 142 GHz feature may be due to blending with the nearby 12–11 line of OC^{34}S . The remaining discrepancy may be explained in terms of our frequency-dependent angular resolution (beam dilution). So far we have assumed that the emission from aziridine has an angular extent on the sky that completely fills the antenna beam pattern at all our observing frequencies. Since hot cores are intrinsically compact objects, this may well not be true. Thus, in a recent study of G327, Gibb et al. conclude that the emission from several of the more complex organics is confined to a region much smaller than the beam sizes listed in Table 2 [20]. A similar situation clearly holds for other similar source regions (e.g. [18]). If the region responsible for our putative aziridine emission is much smaller than the antenna main-beam, the telescope angular resolution (gain) is a factor of 2.5 lower at 142 GHz than at 233 GHz, and a factor of 6.5 lower at 89 GHz than at 233 GHz. Hence, this effect could indeed explain the lack of emission at these lower frequencies. But this also implies that the detected line at 86 312 MHz may be enhanced by blending with emission from an unidentified species.

Table 6
Thiohydroxylamine upper limits^a

Source	T_{mb} (mK: 1σ rms) $3_{1,3}-2_{1,2}$	Linewidth [km s^{-1}]	T_{ex} [K]	$N(\text{NH}_2\text{SH})$ [$\times 10^{13} \text{ cm}^{-2}$]
G327.3–0.6	6.0	5.0	20–50	2.3–4.5
NGC 6334F	11.0	5.0	20–50	4.5–8.2
G10.47+0.03	12.0	5.0	20–50	4.9–8.7
G31.41+0.31	11.0	5.0	20–50	4.5–8.2
G34.3+0.2	11.0	5.0	20–50	4.5–8.2

^a Note: upper limits to the intensity of the indicated transition given in column 2. Upper limits on column density in column 5 are calculated for the estimated line width in column 3 and the range of temperatures in column 4.

Fig. 2.

In summary, the observations are consistent with the assignment of the emission line at 226 233.4 MHz in several sources to the $5_{5,0}-4_{4,1}$ line of aziridine, but further observations are needed to confirm the identification of this molecule in the interstellar medium.

3.2. Propenal (acrolein, CH_2CHCHO)

In its simplest form a rotation diagram analysis plots observed line intensities versus energy of the upper level of each transition, with the intensities normalized by intrinsic line strength in such a way that the plot will be linear if the population distribution is Boltzmann [e.g. [18,58]]. The slope of this line gives the rotational (Boltzmann) temperature, while the intercept is related to the column density. Using such an analysis for the 6 emission lines which coincide with transitions of propenal (Fig. 6), we obtain a reasonable fit to a rotation temperature of 78 ± 20 K in G327.3-0.6 and 96 ± 28 K in SgrB2N, with corresponding column densities of $1.5(10)^{14}$ and $8.7(10)^{14} \text{ cm}^{-2}$, respectively, in the two sources. Using this information, we calculated the expected line strengths for other propenal transitions within our observed spectral regions. These values for T_{rot} and column density predict intensities of the propenal transitions at 88.523, 97.263, and 107.529 GHz (see Table 3 and Table 5) up to an order of magnitude stronger than the 6 lines mentioned above, so that these undetected lines should have been easily observable in our spectra. Although very non-equilibrium intensities are sometimes observed in hot core sources (e.g. stronger than expected emission from weak, b -type transitions, perhaps excited by infrared radiation [18,56]), our failure to detect the propenal transitions with larger A-coefficients leads us to claim only a possible identification of propenal in our sources. We therefore present our results as upper limits to the column density of propenal in Table 7.

Fig. 2. Spectra at 226.2 GHz for several hot core sources (labeled in upper left of each panel), showing the tentative detection of the $5_{5,0}-4_{4,1}$ transition of aziridine. Intensity scale is given as main beam antenna temperature in Kelvin (see text). Other transitions in the band may be assigned to SO_2 , CH_2CHCN , CN, and possibly $c\text{-C}_3\text{HD}$ and ethanol [59,60].

This example of propenal illustrates the importance of observing multiple transitions of a new molecular species, and obtaining a consistent picture of line intensities as a function of expected excitation, before claiming the detection of a new interstellar species. The case of propenal is particularly difficult, because of the large number of transition frequencies which fall into our observed bands.

3.3. Other molecular species

For cyclopropenone, furan, hydroxyethanal, and ethenol we wish to determine an upper limit on the column density along a given line of sight, using the upper limits on the intensity of several transitions. For this purpose we assumed a rather large column density for that species and a uniform excitation temperature (the same for all transitions), and then calculated the corresponding antenna temperatures. Initially such antenna temperatures were higher than our observed limits. We then lowered the column density until the strongest calculated transition had an antenna temperature three times the observed rms noise in the spectrum, at which point all the other transitions had predicted antenna temperatures well below their corresponding noise level (note that this noise level typically differed somewhat among the observed transitions, which may have been observed with different receivers at different times). The procedure was then repeated for another choice of excitation temperature, as indicated in Table 7; the range of excitation temperatures was chosen to match a reasonable range of values for the particular source, as determined from the corresponding excitation of known molecular constituents of that source (see references in Section 1 above). The same procedure, but for only the single observed transition, was followed for NH_2SH (Table 6).

4. Discussion

In comparing the present results with those of previous investigations or with chemical models, it is desirable to estimate the relative abundance,

Fig. 3. Spectra at 86.3 GHz for the hot core sources labeled in upper left of each panel, showing the tentative detection of the $2_{11}-1_{01}$ transition of aziridine ($\text{c-C}_2\text{H}_4\text{NH}$). Intensity scale is antenna temperature in Kelvin (see text). The $J=1-0$ transition of H^{13}CN is in the band.

or limit on the relative abundance, of each species with respect to H_2 , which is by far the most abundant constituent of dense interstellar clouds. As discussed in detail elsewhere [9,18], this is not a simple task. Normally the fractional abundance of species X , symbolized by $[X]$, is approximated by the average along the line of sight given by the ratio of column densities $[X] = \text{N}(X)/\text{N}(\text{H}_2)$. Since the non-polar H_2 is not easily detectable, $\text{N}(\text{H}_2)$ is

usually estimated by assuming an abundance for the principal tracer of cloud material, CO; measuring the column density of one of its rarer isotopomers such as $C^{18}O$ or $C^{17}O$, whose transitions are assumed to be optically thin (unsaturated); and assuming a local isotopic ratio for $^{18}O/^{16}O$ or $^{17}O/^{16}O$.

The values of $N(H_2)$ reported in Table 2 are

Fig. 4. Spectra at 94.4 GHz for several hot core sources (labeled in upper right of each panel), with the frequency of the $21_{1,20}-21_{1,21}$ transition of trans-propenal (C_2H_3CHO) indicated. Intensity scale is given as antenna temperature in Kelvin. Frequency of several symmetry components of the 2-1 transition of 13-carbon methanol are indicated in the top panel.

Fig. 5. Spectra showing 6 transitions of trans-propenal, indicated by their quantum numbers (see Table 3), in the hot core region G327.3-0.6. Intensity scale is given as antenna temperature in Kelvin (see text). Also indicated are transitions of $^{13}CH_3OH$ and cyclopropenylidene- d_1 .

‘beam averages’; that is, they are averages over the angular region defined by the resolution of the telescope at the observed frequency. They have been derived from observations of the CO isotopomer listed in column 7 of Table 2. Some of our observed clouds contain compact, hot, dense cores from which the emission of complex molecules may be enhanced. If such cores are smaller than our angular resolution, we may underestimate the abundance of molecular species

that are present only in such compact regions (cf. [18,20]). Likewise, our beam-averaged upper limits for such species will be lower than the limits which might actually apply to the core itself.

Since we do not have measurements of the CO column densities at the high angular resolutions (of order a few arcsec) which would correspond to such hot core sizes, the abundance upper limits which we shall compute will be beam averages (but see the discussion for aziridine below). Dividing the column density upper limits $N(X)$ from Table 6 and Table 7 by the values of $N(\text{H}_2)$ for each source region from Table 2, we may obtain upper limits for the beam-averaged fractional abundance $[X]$ of each molecular species in each source. These are given in Table 8. Note that we have included propenal in Table 8, because of the uncertainties involving the possible propenal detections described in Section 3.2. In most cases the principal source of uncertainty in the resulting abundances arises from uncertainties in $N(\text{H}_2)$,

which can be as much as an order of magnitude. For comparison with the values in Table 8, we may note that such organic molecules as ethanol ($\text{CH}_3\text{CH}_2\text{OH}$) or methyl formate (CH_3OCHO) have abundances in hot core sources in the range 10^{-9} – 10^{-8} (e.g. [20]).

For aziridine the situation is more complex. Assuming that any aziridine emission would be extended relative to our angular resolution at all observed frequencies (i.e. no beam dilution, consistent with our treatment of the other molecular species described above) and that the energy level populations obeyed a Boltzmann distribution, our failure to detect lines at the frequencies of several aziridine transitions would imply that the lines at 226.2 and 86.3 GHz must be assigned to an unknown carrier(s). The intensity upper limits in Table 4 would then allow us to deduce the column density upper limits given in the second column of Table 9. On the other hand, the 226.2 GHz line can be assigned to the 5_{50} – 4_{41} transition of

Table 7
Column density upper limits^b

Source	$N(\text{CH}_2\text{CHOH})$ [10^{14} cm^{-2}]	$N(\text{c-C}_3\text{H}_2\text{O})$ [10^{12} cm^{-2}]	$N(\text{c-C}_4\text{H}_4\text{O})$ [10^{14} cm^{-2}]	$N(\text{CH}_2\text{OHCHO})$ [10^{12} cm^{-2}]	$N(\text{CH}_2\text{CHCHO})$ [10^{12} cm^{-2}]
TMC-1(CP) ^{a,e}	0.3 ¹ –0.2 ¹	1.7 ⁷ –0.8 ⁷	0.1 ¹¹ –0.2 ¹¹	0.9 ¹⁵ –0.8 ¹⁵	...
Orion KL ^{c,g}	1.9 ² –2.3 ⁶	2.7 ⁹ –4.7 ⁹	2.9 ¹¹ –5.9 ¹⁴	14.0 ¹⁵ –41.1 ¹⁵	...
Orion 3'N ^{b,f}	0.6 ¹ –1.3 ¹	3.6 ⁸ –5.1 ⁸	0.8 ¹¹ –3.0 ¹¹	4.9 ¹⁵ –13.0 ¹⁵	...
G327.3–0.6 ^{c,g}	1.5 ³ –1.2 ⁵	1.3 ¹⁰ –2.1 ¹⁰	1.0 ¹² –1.4 ¹³	...	5.3 ¹⁶ –6.9 ¹⁶
L134N(C) ^{a,e}	0.2 ¹ –0.1 ¹	1.9 ⁸ –0.7 ⁸	0.1 ¹¹ –0.2 ¹¹	0.6 ¹⁵ –0.6 ¹⁵	...
IRS 16293 ^{c,g}	1.3 ² –1.5 ²	8.7 ⁸ –19.0 ⁸	85.0 ¹⁴ –7.1 ¹⁴	...	...
NGC 6334F ^{c,g}	1.1 ³ –1.8 ³	1.3 ¹⁰ –2.1 ¹⁰	1.0 ¹² –1.4 ¹³	...	5.4 ¹⁶ –7.2 ¹⁶
SgrB2 N ^{d,g}	5.8 ³ –6.7 ⁴	2.7 ¹⁰ –4.2 ¹⁰	2.0 ¹² –2.9 ¹³	...	19.0 ¹⁶ –22.0 ¹⁶
G10.47+0.03 ^{c,g}	1.9 ³ –2.3 ⁵	...	...	...	11.0 ¹⁷ –12.0 ¹⁷
W33A ^{c,g}	1.0 ² –1.2 ²	1.6 ⁹ –2.6 ⁹	99.14–9.014	...	...
G31.41+0.31 ^{c,g}	1.9 ³ –2.1 ⁵	...	...	...	11.0 ¹⁷ –12.0 ¹⁷
G34.3+0.2 ^{c,g}	1.9 ³ –2.1 ⁵	...	...	...	...

^a Assuming linewidth of 0.5 km s⁻¹.

^b Assuming linewidth of 3.0 km s⁻¹.

^c Assuming linewidth of 5.0 km s⁻¹.

^d Assuming linewidth of 10.0 km s⁻¹.

^e Assuming excitation temperature range 5 K (first entry) to 10 K (s).

^f Assuming excitation temperature range 10 K (first entry) to 30 K (s).

^g Assuming excitation temperature range 20 K (first entry) to 50 K (s).

^h Note: All values are 3 σ upper limits, although propenal (CH_2CHCHO) may possibly have been detected (see Section 3.2). (1) from transition $2_{1,2}$ – $1_{0,1}$; (2) from transition $4_{2,3}$ – $3_{1,2}$; (3) from transition $5_{1,4}$ – $4_{1,3}$; (4) from transition $6_{3,4}$ – $6_{2,5}$; (5) from transition $8_{1,7}$ – $8_{0,8}$; (6) from transition $12_{1,12}$ – $11_{0,11}$; (7) from transition $5_{1,4}$ – $4_{1,3}$; (8) from transition $6_{0,6}$ – $5_{0,5}$; (9) from transition $7_{1,6}$ – $6_{1,5}$; (10) from transition $8_{0,8}$ – $7_{0,7}$; (11) from transition $2_{0,2}$ – $1_{0,1}$; (12) from transition $9_{0,9}$ – $8_{0,8}$; (13) from transition $11_{0,11}$ – $10_{0,10}$; (14) from transition $22_{0,22}$ – $21_{0,21}$; (15) from transition $4_{1,3}$ – $4_{0,4}$; (16) from transition $10_{0,10}$ – $9_{0,9}$; (17) from transition $11_{0,11}$ – $10_{0,10}$.

Fig. 6. Rotation diagram (see text), showing the 6 apparently detected transitions of *trans*-propenal in the hot core source G327.3 – 0.6. E_u is the energy of the upper level of the transition in temperature units, I is the integrated intensity of the observed line, k is Boltzmann’s constant, μ is the relevant component of the molecular dipole moment, and ν and S are the frequency and intrinsic line strength of the transition (see the discussion of rotation diagrams in [18]). The fit to the data points, which yields the propenal column density N and rotation temperature T_{rot} , ignores the upper limits indicated by the downward-pointing arrows.

aziridine in a consistent manner, if the aziridine emission at 86 GHz is significantly beam-diluted, as would be expected for a hot core. In column three of Table 9 we calculate column densities on the assumption that the aziridine emission comes from a region of the same angular size as our 226 GHz beam size (about 20 arcsec), using $T_{\text{rot}} = 50$ K. The corresponding abundances are listed in column 4. Aziridine abundances in the range $1 - 5 \times 10^{-10}$ with respect to H_2 are deduced. This is about the same as the values found for the abundance of the related species oxirane ($\text{c-C}_2\text{H}_4\text{O}$) in several hot cores [21].

It must be noted that the aziridine column densities in the innermost part of compact hot cores could be considerably higher than the values in Table 9, although the corresponding column densities for H_2 would also be higher. For example, if the emission in the case of the source G327

were confined to a core of angular size about 2 arcsec, as suggested by results for some other large organic species [20], the column density of aziridine would be of order 10^{16} cm $^{-2}$, and the abundance relative to H_2 of order 10^{-9} (the values do not simply scale with the beam area, since the aziridine line becomes saturated). The scaling with rotational temperature goes basically as the partition function, and is thus not a large factor given the other uncertainties.

The case for the tentative detection of interstellar aziridine is stronger than that for propenal for the following reasons. We detect emission lines in several sources at the frequencies of what are expected to be the two strongest aziridine transitions in our observed frequency bands. The failure to detect other aziridine transitions can reasonably be the result of our frequency-dependent angular resolution (beam dilution) towards small,

hot core sources. In contrast, although we detect emission features that might be assignable to six different propenal transitions, we do not see transitions that would reasonably be expected to be considerably stronger. Since all the propenal transitions in question are in the same general frequency range, beam dilution can not explain the discrepancies.

Finally, we should comment on the recent reported detection of hydroxyethanal [73] in the Galactic center cloud SgrB2(N). Since we did not search for hydroxyethanal in this source, no direct comparison is possible. We note, however, that our upper limits in Table 8 for the sources which we did observe are at approximately the same level (a few parts in 10^{10} , using the H_2 column density from Table 2) as the claimed detection in SgrB2(N).

5. Conclusion

We have surveyed a number of interstellar clouds, including both ‘hot core’ sources where massive stars are forming and quiescent cloud regions, for rotational transitions from a number of previously undetected molecular species. We

report the detection of several emission lines that correspond in frequency to transitions of propenal (CH_2CHCHO) and aziridine ($c-C_2H_4NH$). In the case of aziridine, for reasonable source temperatures the 2 detected lines are expected to be the 2 strongest in our observed frequency bands, and we therefore claim the first, tentative, astronomical identification of aziridine. The evidence is strongest in the hot cores G327.3–0.6 and G10.47+0.03. Further observations of other aziridine transitions are needed to confirm this result, which implies abundances of the same order as have been reported for oxirane in similar sources (by number of molecules, a few parts in 10^{10} relative to molecular hydrogen [21]).

For propenal we have only a possible identification. Although some 6 emission features in our spectra of G327.3 and 4 in SgrB2(N) coincide with propenal transitions, other such transitions are not observed. If propenal is indeed responsible for the detected lines, unusual and not understood excitation conditions are present. Observations of intrinsically stronger transitions, at higher frequencies than our present 3 mm data, would be helpful in determining whether propenal is indeed present in interstellar clouds.

Table 8
Abundance upper limits relative to H_2 ^a

Source	CH_2CHOH ($\times 10^{10}$)	$c-C_3H_2O$ ($\times 10^{10}$)	$c-C_4H_4O$ ($\times 10^{10}$)	CH_2OHCHO ($\times 10^{10}$)	CH_2CHCHO ($\times 10^{10}$)	NH_2SH ($\times 10^{10}$)
TMC-1	10	0.8	10	0.5		
Orion KL	20	0.5	60	4		
Orion 3'N	30	1	60	3		
G327.3	5	0.07	5		0.2	2
L134N(C)	9	0.9	10	0.3		
IRS16293	8	1	400			
NGC6334	9	0.1	7		0.4	4
SgrB2 N	2	0.01	1		0.07	
G10.47	20				1	9
W33A	2	0.03	200			
G31.41	10				0.6	4
G34.3	7					3

^a Note: values are 3σ upper limits to the fractional abundance by number, as defined for species X by the ratio of column densities, $N(X)/N(H_2)$; see, however, Section 3.1, regarding the possible detection of CH_2CHCHO . All values are multiplied by 10^{10} . Molecules of similar complexity such as ethanol (CH_3CH_2OH) or methyl formate (CH_3OCHO) have abundances in hot core sources in the range 10^{-9} – 10^{-8} (e.g. [20]).

Table 9

Abundance of aziridine relative to H_2 ^a

Source	$\text{N}(\text{c-C}_2\text{H}_4\text{NH})$ (beam-ave) (10^{13} cm^{-2})	$\text{N}(\text{c-C}_2\text{H}_4\text{NH})$ (small source) (10^{13} cm^{-2})	Abundance ($\times 10^{10}$)
G327.3	<3–6	15	5
NGC6334	<3–6		<3
SgrB2N	≤ 30 –60		1?
G10.47	<6–9	5	5
G31.41	<3–6		<3
G34.3	≤ 3 –6		1?

^a Note: Column 2 lists 3σ upper limit on beam averaged column density, for rotational temperatures of 30 K (left) or 50 K (right). Column 3 lists column density assuming $5_{50-4_{41}}$ line may be assigned to aziridine, implying a source size $\leq 20''$. Column 4 gives the abundance, $\text{N}(\text{c-C}_2\text{H}_4\text{NH})/\text{N}(\text{H}_2)$, multiplied by 10^{10} , based on column 3 if value given, otherwise on column 2.

To our knowledge, there are no published chemical models which include calculations of the expected abundance of aziridine or propenal in interstellar clouds, probably because these species have not hitherto been discussed as potential cloud constituents. Nonetheless, it seems likely that they would be produced by a combination of gas phase reactions and processes in icy grain mantles, such as those discussed in connection with hot cores by Tielens and Charnley [72]. Their calculated abundances, and those of the species for which we obtained only upper limits, would depend, however, on reaction rates and branching ratios for which data is limited or absent, particularly at the temperatures relevant to the interstellar medium. Given simply the relative cosmic abundance of nitrogen to oxygen (about 1:8) and the uncertainties in the measured astronomical abundance of oxirane, the presence of aziridine in hot cores at about our estimated abundance is reasonable.

We also obtained upper limits on the abundance of furan ($\text{c-C}_4\text{H}_4\text{O}$), cyclopropenone ($\text{c-C}_3\text{H}_2\text{O}$), hydroxyethanal (CH_2OHCHO), thiohydroxylamine (NH_2SH), and ethenol (CH_2CHOH) in several regions. Our upper limits for hydroxyethanal are at about the same level as a recently claimed detection of this molecule [73]. All these values will help to constrain the chemical processes that are important in dense interstellar clouds. Future attempts to detect these species, or to improve the present abundance limits, should strive for higher angular resolution toward the small hot core regions. Both the Large Mil-

limeter Telescope, presently being constructed by the University of Massachusetts and the Mexican Instituto Nacional de Astrofisica, Optica y Electronica [65], and millimeter and submillimeter arrays such as CARMA and ALMA [66] will be important in this context. Observations at submillimeter wavelengths would benefit from larger intrinsic line strengths, at least where system sensitivities comparable to those presently available at $\lambda < 1 \text{ mm}$ can be achieved.

Acknowledgements

The authors are grateful to the staffs at SEST, the NRAO 12 m telescope, and the NASA DSN for their assistance and expertise in the observations reported here; to M. Ikeda, E. Gibb, and S.-L. Liu for assistance in carrying out the observations and in reducing the data; to P. Bergman and an anonymous referee for helpful discussions; and to Anne Horn for making measurements of the microwave spectrum of hydroxyethanal. The Swedish-ESO Submillimetre Telescope (SEST) is operated jointly by ESO and the Swedish National Facility for Radio Astronomy, Onsala Space Observatory, at Chalmers University of Technology. The National Radio Astronomy Observatory (NRAO) is a facility of the National Science Foundation, operated under a cooperative agreement by Associated Universities, Inc. This research was supported in part by NASA grants NAG5-7598 (NY Center for Studies of the Origins of Life) and NAG5-8718, by the Swedish

Natural Science Research Council (NFR), by the Deutsche Forschungsgemeinschaft via Grants SFB 301 and SFB 494, and by special funding from the Ministry of Science of the Land Nordrhein-Westfalen.

References

- [1] W.D. Langer, E.F. van Dishoeck, E.A. Bergin, G.A. Blake, A.G.G.M. Tielens, T. Velusamy, D.C.B. Whittet, in: V. Mannings, A.P. Boss, S.S. Russell (Eds.), *Protostars and Planets IV*, University of Arizona Press, Tucson, 2000, p. 29.
- [2] E.F. Van Dishoeck, G.A. Blake, *Ann. Rev. Astron. Astrophys.* 36 (1998) 317.
- [3] Y.C. Minh, E.F. van Dishoeck, *Astrochemistry: From Molecular Clouds to Planetary Systems* (IAU Symp. 197), Astron. Soc. Pacific Conf. Ser., ASP, San Francisco, 2000.
- [4] A.G.G.M. Tielens, C. van Kerckhoven, E. Peeters, S. Hony, in: Y.C. Minh, E.F. van Dishoeck (Eds.), *Astrochemistry: from Molecular Clouds to Planetary Systems* (IAU Symp. 197), Astron. Soc. Pacific Conf. Ser., ASP, San Francisco, 2000, p. 349.
- [5] T.W. Hartquist, D.A. Williams, *The Chemically Controlled Cosmos*, Cambridge University Press, Cambridge, 1995.
- [6] C.D. Ball, M.C. McCarthy, P. Thaddeus, *Astrophys. J.* 528 (2000) L61.
- [7] S. Green, *Ann. Rev. Phys. Chem.* 32 (1981) 103.
- [8] M.B. Bell, J.K.G. Watson, P.A. Feldman, H.E. Matthews, S.C. Madden, W.M. Irvine, *Chem. Phys. Lett.* 136 (1987) 588.
- [9] W.M. Irvine, F.P. Schloerb, Å. Hjalmarson, E. Herbst, in: D.C. Black, M.S. Matthews (Eds.), *Protostars and Planets II*, University of Arizona Press, Tucson, 1985, p. 579.
- [10] E. Herbst, in: W.F. Wall, A. Carraminana, L. Carrasco, P.F. Goldsmith (Eds.), *Millimeter-wave Astronomy: Molecular Chemistry and Physics in Space*, Kluwer, Dordrecht, 1999, p. 341.
- [11] P. Ehrenfreund, W.A. Schutte, in: Y.C. Minh, E.F. van Dishoeck, *Astrochemistry: from Molecular Clouds to Planetary Systems* (IAU Symp. 197), Astron. Soc. Pacific Conf. Ser., ASP, San Francisco, 2000, p. 135.
- [12] J.H. Lacy, J.S. Carr, N.J. Evans, F. Baas, J.M. Achtermann, J.F. Arens, *Astrophys. J.* 376 (1991) 556.
- [13] L.C. Oesterling, S. Albert, F.C. De Lucia, K.V.L.N. Sastry, E. Herbst, *Astrophys. J.* 521 (1999) 255.
- [14] Y. Hoshino, M. Ohishi, K. Akabane, T. Ukai, S. Tsunekawa, K. Takagi, *Astro. Phys.* 104 (1996) 317 Supplementary.
- [15] W.M. Irvine, F.P. Schloerb, J. Crovisier, B. Fegley, M.J. Mumma, in: V. Mannings, A.P. Boss, S.S. Russell (Eds.), *Protostars and Planets IV*, University of Arizona Press, Tucson, 2000, p. 1159.
- [16] S.B. Charnley, M.E. Kress, A.G.G.M. Tielens, T.J. Millar, *Astrophys. J.* 448 (1995) 232.
- [17] T.J. Millar, G.H. Macdonald, A.G. Gibb, *Astron. Astrophys.* 325 (1997) 1163.
- [18] A. Nummelin, P. Bergman, Å. Hjalmarson, P. Friberg, W.M. Irvine, T.J. Millar, M. Ohishi, S. Saito, *Astrophys. J. Suppl.* 128 (2000) 213.
- [19] G.A. Blake, E.C. Sutton, C.R. Masson, T.G. Phillips, *Astrophys. J.* 315 (1986) 621.
- [20] E. Gibb, A. Nummelin, W.M. Irvine, D.C.B. Whittet, P. Bergman, *Astrophys. J.*, (2000) in press.
- [21] A. Nummelin, J.E. Dickens, P. Bergman, Å. Hjalmarson, W.M. Irvine, M. Ikeda, M. Ohishi, *Astron. Astrophys.* 337 (1998) 275.
- [22] L. Olmi, R. Cesaroni, C.M. Walmsley, *Astron. Astrophys.* 307 (1996) 599.
- [23] P. Pratap, J.E. Dickens, R.L. Snell, M.P. Miralles, E.A. Bergin, W.M. Irvine, F.P. Schloerb, *Astrophys. J.* 486 (1997) 862.
- [24] J.E. Dickens, W.M. Irvine, R.L. Snell, E.A. Bergin, F.P. Schloerb, P. Pratap, M.P. Miralles, *Astrophys. J.*, (2000) in press.
- [25] G.A. Blake, E.F. van Dishoeck, D.J. Jansen, T.D. Groesbeck, L.G. Mundy, *Astrophys. J.* 428 (1994) 680.
- [26] C. Ceccarelli, A. Castets, E. Caux, D. Hollenbach, L. Loinard, S. Molinari, A.G.G.M. Tielens, *Astron. Astrophys.* 355 (2000) 1129.
- [27] E. Gibb, D.C.B. Whittet, W.A. Schutte, A.C.A. Boogert, J.E. Chiar, P. Ehrenfreund, P.A. Gerakines, J.V. Keane, A.G.G.M. Tielens, E.F. van Dishoeck, O. Kerkhof, *Astrophys. J.* 536 (2000) 347.
- [28] H. Ungerechts, E.A. Bergin, P.F. Goldsmith, W.M. Irvine, F.P. Schloerb, R.L. Snell, *Astrophys. J.* 482 (1997) 245.
- [29] J.E. Dickens, W.M. Irvine, M. Ohishi, M. Ikeda, S. Ishikawa, A. Nummelin, Å. Hjalmarson, *Astrophys. J.* 489 (1997) 753.
- [30] A. Nummelin, P. Bergman, *Astron. Astrophys.* 341 (1999) L59.
- [31] G. Winnewisser, E. Churchwell, C.M. Walmsley, in: G.W. Chantry (Ed.), *Modern Aspects of Microwave Spectroscopy*, Academic Press, New York, 1979, p. 313.
- [32] M. Winnewisser, G. Winnewisser, T. Honda, E. Hirota, *Z. Naturforsch.* 30 (1975) 1001.
- [33] S. Thorwirth, H.S.P. Müller, G. Winnewisser, *J. Mol. Spec.* 199 (2000) 116.
- [34] W.M. Irvine, M. Senay, A.J. Lovell, H.E. Matthews, D. McGonagle, R. Meier, *Icarus* 143 (1999) 412.
- [35] W.M. Irvine, J.E. Dickens, A.J. Lovell, F.P. Schloerb, M. Senay, E.A. Bergin, D. Jewitt, H.E. Matthews, *Faraday Disc.* 109 (1998) 475.
- [36] F.J. Lovas, R.D. Suenram, W.J. Stevens, *J. Mol. Spec.* 100 (1983) 316.
- [37] H.E. Matthews, W.M. Irvine, *Astrophys. J.* 298 (1985) L61.
- [38] P. Thaddeus, J.M. Vrtilek, C.A. Gottlieb, *Astrophys. J.* 299 (1985) L63.

- [39] S.C. Madden, W.M. Irvine, H.E. Matthews, P. Friberg, D.A. Swade, *Astron. J.* 97 (1989) 1403.
- [40] R.D. Brown, P.D. Godfrey, D.M. Cragg, E.H.N. Rice, W.M. Irvine, P. Friberg, H. Suzuki, M. Ohishi, N. Kaifu, M. Morimoto, *Astrophys. J.* 297 (1985) 302.
- [41] M. Ohishi, H. Suzuki, S.-I. Ishikawa, C. Yamada, H. Kanamori, W.M. Irvine, R.D. Brown, P.D. Godfrey, N. Kaifu, *Astrophys. J.* 380 (1991) L39.
- [42] J.C. Guillemin, G. Wlodarczak, J.C. Lopez, J. Demaison, *J. Mol. Spec.* 140 (1990) 190.
- [43] S. Saito, *Chem. Phys. Lett.* 42 (1976) 399.
- [44] W.M. Irvine, J. Ellde'r, Å. Hjalmarson, E. Kollberg, O.E.H. Rydbeck, G.O. Sørensen, B. Bak, H. Svanholt, *Astron. Astrophys.* 97 (1981) 192.
- [45] P.C. Myers, P. Thaddeus, R.A. Linke, *Astrophys. J.* 241 (1980) 155.
- [46] M.L. Kutner, D.E. Machnik, K.D. Tucker, R.L. Dickman, *Astrophys. J.* 242 (1980) 541.
- [47] G. Wlodarczak, L. Martinache, J. Demaison, *J. Mol. Spec.* 127 (1988) 200.
- [48] R. Krishnamurthy, G. Arrhenius, A. Eschenmoser, *Origins Life Evol. Biosphere* 29 (1999) 333.
- [49] K.-M. Marstokk, H. Møllendal, *J. Molec. Structure* 5 (1970) 205.
- [50] W.M. Irvine, *Space Sci. Revs.* 90 (1999) 203.
- [51] P. Bergman, Tech. Rpt. 227, School Elec. Computer Eng., Chalmers Univ. Technology, Gothenburg, Sweden, 1992.
- [52] F.F.S. van der Tak, E.F. van Dishoeck, N.J. Evans, G.A. Blake, *Astrophys. J.* 537 (2000) 283.
- [53] W.M. Irvine, *Origins Life Evol. Biosphere* 28 (1998) 365.
- [54] J. Ellde'r, P. Friberg, Å. Hjalmarson, B. Höglund, W.M. Irvine, L.E.B. Johansson, H. Olofsson, G. Rydbeck, O.E.H. Rydbeck, *Astrophys. J.* 242 (1980) L93.
- [55] D.M. Mehringer, L.E. Snyder, Y. Miao, F.J. Lovas, *Astrophys. J.* 480 (1997) L71.
- [56] B.E. Turner, *Astrophys. J.* 76 (1991) 617 Supplementary.
- [57] W.M. Irvine, P.F. Goldsmith, Å. Hjalmarson, In: D.J. Hollenbach, H.A. Thronson, *Interstellar Processes*, D. Reidel, Dordrecht, 1987, p. 561.
- [58] L.E.B. Johansson, C. Andersson, J. Ellde'r, P. Friberg, Å. Hjalmarson, B. Höglund, W.M. Irvine, H. Olofsson, G. Rydbeck, *Astron. Astrophys.* 130 (1984) 227.
- [59] F.J. Lovas, *J. Phys. Chem. Ref. Data* 21 (1992) 181.
- [60] H.M. Pickett, R.L. Poynter, E.A. Cohen, M.L. Delitsky, J.C. Pearson, H.S.P. Muller, *J. Quant. Spectrosc. Rad. Transfer* 60 (1998) 883 <http://spec.jpl.nasa.gov/>.
- [61] J. Cernicharo, In: Y.C. Minh and E.F. van Dishoeck, *Astrochemistry: from Molecular Clouds to Planetary Systems* (IAU Symp. 197), Astron. Soc. Pacific Conf. Ser., ASP, San Francisco, 2000, p. 375.
- [62] H. Feuchtgruber, F.P. Helmich, E.F. van Dishoeck, C.M. Wright, *Astrophys. J.* 535 (2000) L111.
- [63] S. Thorwirth, R. Gendriesch, H.S.P. Müller, F. Lewen, G. Winnewisse, *J. Mol. Spec.* 201 (2000) 323.
- [64] J. Binney, M. Merrifield, *Galactic Astronomy*, Princeton University Press, Princeton, NJ, 1998.
- [65] J.W.M. Baars, L. Carrasco, F.P. Schloerb, *Bull. Amer. Astron. Soc.* 31 (1999) 09–15.
- [66] J. Mangum, *Imaging at Radio through Submillimeter Wavelengths*, Astron. Soc. Pacific, San Francisco, 2000.
- [67] B.P. van Eijck, E. van Zoonen, *J. Mol. Spectrosc.* 111 (1985) 138.
- [68] A. Bauder, *J. Phys. Chem. Ref. Data* 8 (1979) 583.
- [69] (a) T.H. Dunning, *J. Chem. Phys.*, 90 (1989) 1007; (b) R.A. Kendall, T.H. Dunning, R.J. Harrison, *J. Chem. Phys.*, 96 (1992) 6796; (c) T.H. Dunning, D.E. Woon, *J. Chem. Phys.*, 98 (1993) 1358; (d) T.H. Dunning, D.E. Woon, *J. Chem. Phys.*, 100 (1994) 2975; (e) T.H. Dunning, D.E. Woon, *J. Chem. Phys.*, 103 (1995) 4572.
- [70] C. Møller, M.S. Plesset, *Phys. Rev.* 46 (1934) 618.
- [71] Gaussian 98, Revision A.7, M.J. Frisch, G.W. Trucks, H.B. Schlegel, G.E. Scuseria, M.A. Robb, J.R. Cheeseman, V.G. Zakrzewski, J.A. Montgomery, Jr., R.E. Stratmann, J.C. Burant, S. Dapprich, J.M. Millam, A.D. Daniels, K.N. Kudin, M.C. Strain, O. Farkas, J. Tomasi, V. Barone, M. Cossi, R. Cammi, B. Mennucci, C. Pomelli, C. Adamo, S. Clifford, J. Ochterski, G.A. Petersson, P.Y. Ayala, Q. Cui, K. Morokuma, D.K. Malick, A.D. Rabuck, K. Raghavachari, J.B. Foresman, J. Cioslowski, J.V. Ortiz, A.G. Baboul, B.B. Stefanov, G. Liu, A. Liashenko, P. Piskorz, I. Komaromi, R. Gomperts, R.L. Martin, D.J. Fox, T. Keith, M.A. Al-Laham, C.Y. Peng, A. Nanayakkara, C. Gonzalez, M. Challacombe, P.M.W. Gill, B. Johnson, W. Chen, M.W. Wong, J.L. Andres, C. Gonzalez, M. Head-Gordon, E.S. Replogle, J.A. Pople, Gaussian, Inc., Pittsburgh PA, 1998.
- [72] A.G.G.M. Tielens, S.B. Charnley, *Origins Life Evol. Biosphere* 27 (1997) 23.
- [73] J.M. Hollis, F.J. Lovas, P.R. Jewell, *Astrophys. J.* (2000) in press.
- [74] D.C.B. Whittet, *Dust in the Galactic Environment*, Institute of Physics Publishing, Bristol, 1992.